arXiv:cond-mat/0610659v2 [cond-mat.mes-hall] 13 Feb 2007

Oscillating Casimir force between impurities in one-dimensional Fermi liquids

J.N. Fuchs,[1] A. Recati,[2] and W. Zwerger[3]

[1] *Laboratoire de Physique des Solides, Université Paris-Sud, CNRS, UMR 8502, F-91405 Orsay, France.*
[2] *CRS BEC-INFM, Povo and ECT*, Villazzano, I-38050 Trento, Italy.*
[3] *Technische Universität München, James Franck Strasse, D-85748 Garching, Germany.*

(Dated: February 6, 2008)

We study the interaction of two localized impurities in a repulsive one-dimensional Fermi liquid via bosonization. In a previous paper [Phys. Rev. A **72**, 023616 (2005)], it was shown that at distances much larger than the interparticle spacing the impurities interact through a Casimir-type force mediated by the zero sound phonons of the underlying quantum liquid. Here we extend these results and show that the strength and sign of this Casimir interaction depend sensitively on the impurities separation. These oscillations in the Casimir interaction have the same period as Friedel oscillations. Their maxima correspond to tunneling resonances tuned by the impurities separation.

Interacting one-dimensional fermions are nowadays available in many experimental systems. They can be electrons in semiconductor quantum wires, in carbon nanotubes, in conducting polymers, etc. [1] or fermionic atoms in ultra-cold gases [2]. Impurities have a profound effect on the physical properties of these low-dimensional systems. In certain exceptionally clean systems, impurities can be created and controlled. For example, it was recently proposed that atomic quantum dots – i.e. individually trapped atoms playing the role of impurities – could be created in an ultra-cold atomic gas [3]. Another possibility rests in the "buckles" artificially created with AFM tips in single-wall carbon nanotubes and which behave as tunnel junctions [4]. As a last example, we mention atom wires that were recently created on a Si(553)-Au surface and which featured vacancy-like defects that were manipulated with an STM tip [5]. Once it is recognized that impurities can be manipulated, one can ask new questions in which they appear as main actors. For example, what is the interaction between impurities which is mediated by the quantum liquid of fermions? This problem was recently addressed in [6]. It was found that when the impurities are far apart they behave like acoustic mirrors and interact via a Casimir-like force. This is the expected behavior for any system whose low energy properties are described in terms of phonons. However, for the special case of non-interacting fermions, the calculation can be done exactly and it was shown that this interaction features oscillations as a function of the impurities separation [6]. In the present paper – part II – which can be considered as a continuation of [6] – to which we shall refer as part I – we extend some of these results to interacting fermions. Most significantly, we show that the oscillation phenomenon in the interaction remains essentially unaffected in the case of interacting (repulsive) fermions. In the light of proofs showing the absence of repulsive Casimir interactions for the photonic field [7], this is a quite remarkable situation. Here we have a Casimir force whose strength and sign can be tuned by the impurities separation.

In order to investigate that problem, we consider the following model: two localized delta impurities embedded in a repulsive one-dimensional Fermi gas, whose low-energy behavior is that of a single channel Luttinger liquid with parameter $K \leq 1$. The interaction between the impurities – which is mediated by the quantum fluctuations in the liquid – is considered in the adiabatic limit in which changing the distance between the impurities is slow compared to the time the system takes to equilibrate. Within such a limit, the interaction energy directly follows from the groundstate energy. We should stress that we focus on interactions mediated by quantum zero-point fluctuations and not on classical effects due to modification of the average particle density. The latter dominate at short distances of the order of the inter-particle distance and lead to a kind of "mattress effect". For a discussion of the classical ground state energy, see part I.

We first consider non-interacting fermions and discuss the oscillations in the interaction energy in that case. Two delta impurities located at $x_1 = 0$ and $x_2 = r$ are embedded in a one dimensional (1D) Fermi gas with Fermi velocity $v_F = \pi\rho_0/m$ where $\rho_0 = k_F/\pi$ is the average density. The impurities interact with the particles through a potential $g_\alpha \delta(x - x_\alpha)$ where $\alpha = \{1, 2\}$ and g_α are microscopic scattering amplitudes. When the distance between the impurities r is much larger than the average inter-particle distance $1/\rho_0$, it was shown in part I that the renormalized interaction energy between the two impurities is given by

$$V_{12} = \frac{v_F}{2\pi r} \Re \, \mathrm{Li}_2 \big[-\frac{\gamma_1\gamma_2}{\sqrt{1+\gamma_1^2+\gamma_2^2+(\gamma_1\gamma_2)^2}} e^{i\chi} \big] \, . \quad (1)$$

In the preceding equation, Li_2 is the di-logarithmic function, $\Re$ denotes the real part, $\gamma_\alpha = g_\alpha/v_F$ is the dimensionless impurity strength and we introduced the phase $\chi(r, \gamma_\alpha)$ such that $\chi \equiv 2k_F r + \delta$ modulo 2π and $-\pi < \chi \leq \pi$. The phase shift $\delta(\gamma_\alpha) = -\arctan\gamma_1 - \arctan\gamma_2 = -\arctan[(\gamma_1 + \gamma_2)/(1 - \gamma_1\gamma_2)]$ reflects the time delay of fermions scattering on delta impurities. As the function $V_{12}/(v_F/r)$ is 2π-periodic in χ, the interaction energy V_{12} oscillates as a function of r. Maxima are separated by $\Delta(2k_F r) = 2\pi$, which corresponds to the average inter-particle distance $\Delta r = 1/\rho_0$, just as Friedel oscillations. Maxima occur when $\cos(\chi/2) = \cos(k_F r + \delta/2) = 0$.



Figure 1: Interaction energy V_{12} between two identical strong impurities separated by a distance r. Dashed line: non-interacting fermions $V_{12} = v_F \Re\,\mathrm{Li}_2(-e^{i\chi})/2\pi r$. Full line: interacting fermions (with Luttinger parameter $K = 0.6$) $V_{12} = -\pi v_F/24Kr + v_F\chi^2/8\pi r K^2$. In both cases, $\chi = 2k_F r + \delta$. For non-interacting fermions and strong impurities ($\gamma \gg 1$) the phase shift $\delta = -2\arctan\gamma \approx -\pi$. For interacting fermions ($K < 1$) and strong impurities ($\widetilde{\gamma} \gg 1$), we expect the phase shift to be $\delta = -2\arctan(K^2\widetilde{\gamma}) \approx -\pi$, see the main text. The full line corresponds to this guess $\delta \approx -\pi$.

This is precisely the condition for tunneling resonances in a non-interacting 1D Fermi gas, as shown in Appendix A. In Appendix B, we discuss an interesting situation in which the oscillations can disappear.

In the limit of strong impurities ($\gamma_\alpha \gg 1$), the phase shift $\delta \approx -\pi$ and Eq. (1) becomes

$$V_{12} \approx \frac{v_F}{2\pi r}\Re\,\mathrm{Li}_2(-e^{i\chi}) = -\frac{\pi v_F}{24r} + \frac{v_F\chi^2}{8\pi r}\,, \qquad (2)$$

where $\chi \approx 2k_F r - \pi$. The last equation applies to $-\pi < \chi \leq \pi$ with a periodic extension beyond that. The first term has exactly the form of a 1D Casimir energy $-\pi v_F/24r$, with the Fermi velocity v_F playing the role of light velocity, see e.g. [8]. The second term introduces oscillations. Indeed, the function $V_{12}/(v_F/r)$ is 2π-periodic in χ and features wide maxima made of pieces of parabolas and centered at $\chi = \pi$, see Figure 1. On the one hand, at a minimum, where $\chi = 0$, the interaction energy is given by the Casimir energy $-\pi v_F/24r$. This correspond to a situation of no resonant tunneling (anti-resonances). The Casimir energy has the usual interpretation in terms of phononic "radiation pressure" on the impurities: the outer radiation pressure of phonons overcomes the inner one and hence the resulting attraction between the impurities. On the other hand, at a maximum, where $\chi = \pi$, the interaction energy changes sign and becomes positive reaching $\pi v_F/12r$. A physical understanding of this change of sign in the Casimir force might be obtained by noting that in a situation of resonant tunneling ($\chi = \pi$), fermions approaching the double barrier structure from outside can tunnel with no reflection. There is thus no contribution to the radiation pressure due to the phonons in the region outside the impurities. By contrast, fermions which come from in between the two barriers, have to tunnel through a single impurity and do not benefit from resonant tunneling. At low energies, they are almost perfectly reflected and therefore contribute a positive radiation pressure on the impurities. At resonance therefore, the inner radiation pressure is larger than the outer one, and thus the effective interaction between the impurities is repulsive. In a sense, this is like having impurities with transmission properties that are not right-left symmetric and makes a connection to Imry's idea of tuning the sign of the QED Casimir effect using asymmetric mirrors [9].

We now come to the main point of part II. We show that the basic oscillation phenomenon contained in Eq (2), and found in part I for non-interacting fermions, remains essentially intact when including interactions between the fermions. In order to study 1D interacting fermions, we use the Luttinger liquid formalism [10] and concentrate on repulsive interactions. For simplicity, we consider spinless fermions, for which the low-energy description is given by the following hydrodynamic action

$$S_0 = \frac{1}{2\pi K}\int \mathrm{d}x \int_0^\beta \mathrm{d}\tau \left[u(\partial_x\theta)^2 + \frac{1}{u}(\partial_\tau\theta)^2\right]\,, \qquad (3)$$

where $\beta = 1/T$ is the inverse temperature and θ is related to the density of the liquid by

$$\rho(x) = (\rho_0 + \partial_x\theta/\pi)\,[1 + 2\cos(2\theta + 2k_F x) + \ldots]\,, \quad (4)$$

where ρ_0 is the equilibrium density and only the first harmonics are written [10]. The Luttinger liquid description depends on three parameters: the sound velocity u, the Luttinger interaction parameter K and a cutoff energy $\omega_c \sim uk_F$ above which this effective description breaks down. Repulsive short-range interactions imply $0 < K < 1$. With long-range (Coulomb) interactions, the system becomes a 1D Wigner crystal, which can be seen as the $K \to 0$ limit of the Luttinger liquid [11, 12], but can not be treated within the present formalism. Non-interacting fermions correspond to $K = 1$ and $u = v_F$. If there is full translational invariance $uK = v_F$ [10].

We consider, as before, two localized impurities sitting at position $x_1 = 0$ and $x_2 = r$. The interaction between the impurities and the Luttinger liquid is

$$S_i = \int_0^\beta \mathrm{d}\tau \sum_{\alpha=1,2} \widetilde{g}_\alpha \rho(x_\alpha)\,, \qquad (5)$$

where $\widetilde{g}_\alpha$ is a phenomenological scattering amplitude. As the Luttinger liquid is an effective description valid at low energy, the phenomenological scattering amplitude $\widetilde{g}_\alpha$ is not equal to the microscopic scattering amplitude g_α. The precise relation between the two scattering amplitudes is not generally known. In the renormalization group language, the phenomenological scattering amplitude $\widetilde{g}$ is roughly a running scattering amplitude at the energy scale ω_c. For simplicity, we consider the symmetric case $\widetilde{g}_1 = \widetilde{g}_2 = \widetilde{g}$. Inserting the density expansion (4) into the interaction (5) gives rise to several different terms. The first term $2\beta\widetilde{g}\rho_0$ is just twice the classical energy cost of adding an impurity to the liquid: it does not contribute to the renormalized interaction energy V_{12}. The second term due to forward scattering is proportional to $\partial_x\theta$. This term is often eliminated via a gauge transformation: $\theta(x) \to \theta(x) - K\widetilde{g}(\mathcal{H}(x-r) + \mathcal{H}(x) - 1/2)/u$, where $\mathcal{H}(x)$ is the Heaviside step function. However, this operation is not totally innocent here and forward scattering can not be eliminated: the gauge transformation gives rise to a phase shift $\delta = -2K\widetilde{g}/u$ in the $2k_F$ backscattering term. In the limit of weak impurities, this phase shift agrees with the one found for non-interacting fermions $\delta = -2\arctan(g/v_F) \approx -2g/v_F$. Actually, we expect that the phase shift in the interacting case is given by $\delta = -2\arctan(K\widetilde{g}/u)$ for *arbitrary* impurity strengths, however to our knowledge there are no exact results on this problem. In fact, the precise value of δ depends on the detailed potential and the behavior of the fermion liquid at high energies where a LL description no longer applies. This issue is often discussed in the context of the x-ray edge singularity or in the Kondo problem and is related to the order in which two non-commuting limits are taken: sending the size of the impurity to zero versus sending the bandwidth to infinity. When using the Luttinger liquid formalism, one automatically takes the second limit first, see Ref. [13]. The third term due to $2k_F$ backward scattering gives the dominant contribution to the action S_i and, when including the phase shift δ, appears as

$$S_i \approx V\int_0^\beta \mathrm{d}\tau(\cos[2\theta(0)] + \cos[2\theta(r)+\chi]) \;, \qquad (6)$$

where $V = 2\widetilde{g}\rho_0$ and χ is a phase such that $\chi \equiv 2k_F r + \delta$ modulo 2π and $-\pi < \chi \le \pi$. There are also higher order terms coming from the density harmonics, however their naive scaling dimension shows that they are less relevant and we therefore neglect them. Actually perturbative renormalization group calculations show that $2k_F$ backscattering is always relevant as long as $K<1$ while all higher backscattering operators $\cos(2n\theta(x) + 2nk_F x)$, with $n = 2, 3, \ldots$, are irrelevant as long as $K > 1/n^2$ [14, 15]. As we neglect those higher backscattering operators, our model only describes the situation of two impurities in an interacting Fermi gas such that $1/4 < K < 1$.

As a side remark, we note that there are other ways of coupling impurities to a bosonic bath of oscillators, such as a Luttinger liquid. For example, in the context of QED Casimir force studies [16], the impurities (mirrors) are usually taken to couple directly to the density of the photonic field, i.e. θ^2 rather than $\cos 2\theta$ would appear in equation (6). This is the case in particular for the theorem of Kenneth and Klich on the absence of a repulsive Casimir force [7]. Another example was considered in Ref. [3], where only forward scattering was kept, resulting in a mapping to the spin-boson model. These different models, which all describe two impurities in a bath of quantum oscillators, do not necessarily show the same behavior.

From now on, we shall work with the action $S = S_0 + S_i$ with S_i given by Eq. (6). The partition function is $Z = \int \mathcal{D}\theta(x,\tau)\exp(-S)$ and the integral on the field $\theta(x,\tau)$ can be performed everywhere except at the position of the impurities $x = x_\alpha$. In this way we are left with an action $A = A_0 + A_i$ for the fields $\theta_\alpha(\tau) \equiv \theta(x_\alpha, \tau)$ only

$$\begin{aligned} A_0 &= \sum_{n\in\mathbb{Z}} \frac{f_n}{\pi K}(|\theta_{1,n}|^2 + |\theta_{2,n}|^2 - e_n(\theta_{1,-n}\theta_{2,n} + \text{c.c.})) \\ A_i &= V\int_0^\beta \mathrm{d}\tau(\cos[2\theta_1] + \cos[2\theta_2+\chi])\,, \qquad (7) \end{aligned}$$

where ω_n are bosonic Matsubara frequencies, the Fourier transform is defined as $\theta_n = T\int d\tau \exp(i\omega_n\tau)\theta(\tau)$, the characteristic finite-size frequency is $\omega_r = u/r$ and we defined $e_n = \exp(-|\omega_n|/\omega_r)$ and $f_n = \beta|\omega_n|/(1-e_n^2)$. Up to an overall normalization factor, the partition function is now given by $Z = \int \mathcal{D}\theta_\alpha(\tau)\exp(-A)$. The partition function gives the free energy $F = -T\ln Z$, from which the renormalized interaction energy follows: $V_{12}(r) = F(r) - F(\infty)$. From the parametric dependance of the action A on the phase χ, we see that the free energy (and therefore V_{12}) will be 2π-periodic in χ. As $\chi = 2k_F r + \delta$, this means that V_{12} will have some structure with periodicity $1/\rho_0$ in r just as for non-interacting fermions. Therefore, the interaction energy V_{12} oscillates as a function of the impurities separation r.

We first evaluate the partition function using the saddle point approximation. For small K and strong impurities $\widetilde{\gamma} = \widetilde{g}/v_F \gg 1$, the fields θ_α are pinned to the values that minimize A_i. We therefore expand the cosines around a minimum to quadratic order in the fields. It should be noted that terms independent of the fields give an important contribution here [20]. The resulting free energy is:

$$\begin{aligned} F_s(V) &= F_s(0) + \frac{\omega_r\chi^2}{8\pi K}\frac{2\pi KV}{\omega_r + 2\pi KV} \qquad (8) \\ &+ \frac{T}{2}\sum_n \ln(1 + \frac{2\pi KV}{|\omega_n|}(1+e_n))(1 + \frac{2\pi KV}{|\omega_n|}(1-e_n)) \end{aligned}$$

As the validity of the saddle point approximation requires that $2\pi KV \gg \omega_r$, the renormalized interaction becomes:

$$\begin{aligned} V_{12} &= \frac{\omega_r\chi^2}{8\pi K} \qquad (9) \\ &+ \frac{T}{2}\sum_n \ln\frac{1+\frac{2\pi KV}{|\omega_n|}(1+e_n)}{1+\frac{2\pi KV}{|\omega_n|}}\frac{1+\frac{2\pi KV}{|\omega_n|}(1-e_n)}{1+\frac{2\pi KV}{|\omega_n|}} \end{aligned}$$

In the limit of zero temperature $\omega_r \gg T$, it gives

$$\begin{aligned} V_{12} - \frac{\omega_r\chi^2}{8\pi K} &\approx \frac{\omega_r}{2\pi}\int_0^{\omega_c/\omega_r} dx\ln(1-e^{-2x}) \\ &\approx -\frac{\pi\omega_r}{24}\,, \qquad (10) \end{aligned}$$

where we used $\omega_c \gg \omega_r$ [21]. This result generalizes Eq. (2) when interactions between particles are taken into account. It also admits an interpretation as an attractive Casimir energy $-\pi\omega_r/24$ – albeit with the true sound velocity u replacing the Fermi velocity v_F – modified by oscillations contained in the term $\omega_r\chi^2/8\pi K$, see Figure 1. The width of the maxima measured relative to their amplitude is proportional to K. Therefore the maxima are narrower in the presence of interactions between the particles. The exact position of the maxima is given by the values of r such that $\chi = \pi$, i.e. $\cos(\chi/2) = \cos(k_F r + \delta/2) = 0$, and depends on both the impurity strength and the inter-particle interactions through the phase shift δ. As in the case of non-interacting fermions, this is exactly the condition at which tunneling resonances occur in a Luttinger liquid, see the Appendix A.

At this stage, it is worth mentioning that there is a mapping of our problem – as given by actions (3) and (6) – on the double-boundary sine-Gordon model studied in the context of conformal field theory [17]. For special values of the Luttinger parameter $K = 1/[2(1+n)]$, where n is a strictly positive integer (the so-called reflectionless points), the authors of Ref. [17] were able to calculate exactly the groundstate energy of this model [22]. In the limit of strong impurities and for the reflectionless points, it yields the same interaction energy $V_{12} = -\pi\omega_r/24 + \omega_r\chi^2(1+n)/4\pi$ as Eq. (10).

In order to study what happens for values of K close to one, we turn to the self-consistent harmonic approximation (SCHA), which is just an implementation of Feynman's variational principle, see, e.g., [1]. Expanding the cosines in Eq. (6) to second order in the fields, we replace the coefficient $2V$ in front of the quadratic term by a variational parameter λ. The true action $S = S_0 + S_i$ is now replaced by a quadratic variational action S_v. The parameter λ is determined self-consistently by finding a minimum to the variational free energy $F_v(\lambda) = -T\ln Z_v + T\langle S - S_v\rangle$. The latter is given by

$$F_v(\lambda) = F_s(0) + f - \lambda\frac{\partial f}{\partial\lambda} - 2V\exp(-\frac{\partial f}{\partial\lambda}) \tag{11}$$

where

$$\begin{aligned} f &= F_s(V=\lambda/2) - F_s(0) \\ &\approx -\frac{\pi\omega_r}{24} + \frac{\omega_r\chi^2}{8\pi K} + \frac{\omega_c}{\pi}\ln(1+\frac{\pi K\lambda}{\omega_c}) + K\lambda\ln(1+\frac{\omega_c}{\pi K\lambda}) \end{aligned}$$

the second equality being justified when $\omega_c \gg \omega_r \gg T$ and $\pi K\lambda \gg \omega_r$ which is a validity condition of the SCHA. Extremizing the variational free energy gives:

$$\lambda = 2V\exp(-\frac{\partial f}{\partial\lambda}) = 2V\left(\frac{\pi K\lambda}{\pi K\lambda + \omega_c}\right)^K \tag{12}$$

When $K < 1$, this equation has two solutions: 0 (corresponding to a maximum of F_v) and $\bar\lambda > 0$ (corresponding to a minimum), where $\bar\lambda \approx 2V$ when $2\pi V \gg \omega_c$ and $\bar\lambda \approx 2V(2\pi KV/\omega_c)^{K/(1-K)}$ when $2\pi V \ll \omega_c$ [23]. As neither $\bar\lambda$ nor $\partial f(\bar\lambda)/\partial\lambda$ depends on r, the renormalized interaction energy is given by $V_{12} \approx f(\bar\lambda, r) - f(\bar\lambda, \infty) \approx -\pi\omega_r/24 + \omega_r\chi^2/8\pi K$. The result obtained in the saddle point approximation Eq. (10) is recovered and is therefore valid more generally for $K < 1$ as long as $\pi K\bar\lambda \gg \omega_r$, i.e. for strong impurities or large distances between the impurities.

The situation of weak impurities or short distances (but always in the limit $r \gg 1/\rho_0$) can be studied thanks to a perturbative calculation of the partition function. In the regime $K > 1/2$, the interaction energy is

$$V_{12} = -\tilde\gamma^2\frac{v_F}{2\pi r}\cos\chi \times \frac{(uk_F)^2 K\Gamma(K-1/2)}{\omega_c^{2K}\omega_r^{2-2K}\sqrt{\pi}\Gamma(K)}\ , \tag{13}$$

where $\Gamma(x)$ is the gamma function and $\chi \approx 2k_F r$ when $\tilde\gamma \ll 1$, as already shown in part I. This is a RKKY-type interaction, which decays with a renormalized power law r^{1-2K}. It is valid as long as $|V_{12}| \ll \omega_r$, i.e. $r \ll \rho_0^{-1}\tilde\gamma^{1/(K-1)}$. In a similar way, we obtain

$$V_{12} = -\tilde\gamma^2\cos\chi \times \frac{2(v_F k_F)^2\beta}{\pi^2(1-2K)(1-K)(\beta\omega_c)^{2K}}\ , \tag{14}$$

in the regime $0 \le K < 1/2$ and

$$V_{12} = -\tilde\gamma^2\cos\chi \times \frac{4(v_F k_F)^2}{\pi^2\omega_c}\ln(\beta\omega_r)\,, \tag{15}$$

for the special case $K = 1/2$, again provided that $\omega_r \gg T$. For $0 \le K < 1/2$, the interaction given by Eq. (14) still shows Friedel oscillations but its envelope does not depend on the inter-impurity distance r anymore, which reflects the tendency of the system toward Wigner crystallization when K goes to zero. However the validity of this calculation is restricted to small but finite temperature as $|V_{12}| \ll \omega_r$ corresponds to $r \ll u\omega_c^{2K}T^{1-2K}/(v_F k_F\tilde\gamma)^2$. This means that when $T \to 0$, the domain of validity of perturbation theory shrinks to zero, when $0 \le K \le 1/2$. We note that, at the reflectionless point $K = 1/4$, Eq. (14) has the same structure as the exact $T = 0$ result $V_{12} = -\tilde\gamma^2\cos\chi \times (v_F k_F)^2/(\pi\omega_c)$ [17], where $\chi \approx 2k_F r$ when $\tilde\gamma \ll 1$. Again, as in the case of strong impurities, maxima in the interaction energy V_{12} occur when $\cos(\chi/2) = 0$, which is the condition for tunneling resonances in a Luttinger liquid, see the Appendix A.

Tunneling resonances for the conductance are only found for $1/4 < K \le 1$. Indeed on resonance $2k_F$ backscattering is absent and higher order backscattering operators control the behavior of the system. Once $K < 1/4$, $4k_F$ backscattering, which we neglected, becomes relevant and washes out the tunneling resonances [14]. Therefore, in order to study what happens for very small values of K, one should also take these other backscattering operators into account when computing V_{12}. Using the methods of part I, the present work could also be extended to treat spin 1/2 fermions.

In conclusion, we found that at large impurities separation or for strong impurities, the renormalized interaction between the impurities results from a Casimir force modified by oscillations – due to quasi-stationary states in between the impurities. The oscillations can be spanned by changing the distance between the impurities and are separated by the average inter-particle distance, i.e. they have the same periodicity as Friedel oscillations. The position of the maxima (peaks) is given by precisely the same condition as for tunneling resonances and depends on particle interactions – through the Luttinger parameter K – and on the impurity strength through a phase shift. At intermediate distances and for Luttinger parameter $K > 1/2$, the interaction is of the RKKY type, featuring Friedel oscillations with an envelope which is decaying as a power law with a renormalized exponent. When $1/4 < K < 1/2$, Friedel oscillations are still present at finite T but are not decaying anymore. With ultracold fermionic atoms confined in a cigar-shaped trap, as described in detail in part I, it should be possible to detect this oscillating Casimir force.

Acknowledgments

We acknowledge useful discussions with I. Safi, H. Saleur, N. Prokof'ev and P. Faccioli. We also thank V. Meden for very instructive discussions and comments on the present manuscript.

Appendix A: TUNNELING RESONANCES

We first consider a single fermion at the Fermi surface – with incoming wave vector k_F – scattering on two delta impurities $g_1\delta(x)$ and $g_2\delta(x-r)$. Using the transport matrix formalism, it is easy to obtain its transmission probability across the double barrier structure

$$\begin{aligned} P(k_F) &= \Big[1+\gamma_1^2+\gamma_2^2+2\gamma_1^2\gamma_2^2 \\ &+ 2\gamma_1\gamma_2\sqrt{(1+\gamma_1^2)(1+\gamma_2^2)}\cos(2k_F r+\delta)\Big]^{-1}, \end{aligned} \tag{A1}$$

where $\gamma_j = mg_j/k_F$ and $\delta = -\arctan\gamma_1 - \arctan\gamma_2$. This probability has maxima – called tunneling resonances – when $\cos(k_F r+\delta/2)=0$. On resonance, the energy of a quasi-stationary state in between the impurities matches the Fermi energy. In the case of symmetric impurities ($\gamma_1=\gamma_2$), there is perfect transmission on resonance. For asymmetric impurities, the transmission probability is always smaller than one. The condition for tunneling resonances coincides with the position of maxima of V_{12}.

We now turn to interacting fermions. Tunneling resonances in a Luttinger liquid were first studied in Ref. [14, 15] and are reviewed by Giamarchi [18]. In these works, the forward scattering term on the impurities was not taken into account. When this term is neglected, and in the absence of a gate voltage, the criterion for tunneling resonances in a Luttinger liquid is $\cos(k_F r)=0$ [14, 15, 18]. It is trivial to redo these calculations starting from the action (6) including the phase shift $\delta = -2K\widetilde{g}/u$ resulting from forward scattering on the impurities. The only modification amounts to replace $2k_F r$ by $2k_F r+\delta$ and therefore the criterion for tunneling resonances becomes $\cos(k_F r+\delta/2)=0$. As in the case of non-interacting fermions, this coincides with the position of the maxima in V_{12}.

Appendix B: DISAPPEARANCE OF THE OSCILLATIONS IN V_{12}

Here we discuss an interesting phenomenon in the interaction energy V_{12}, which was pointed out to us by V. Meden [19]. Consider non-interacting spinless fermions on a lattice in a tight-binding model with hopping amplitude t, lattice spacing b, and filling factor $f=\rho_0 b = 1/2$. The Fermi momentum is $k_F=\pi f/b = \pi/(2b)$ and the Fermi velocity is $v_F = 2tb\sin(f\pi)=2tb$. This model is expected to have the same low energy behavior as the continuum model considered in the present paper. There are also two site impurities with dimensionless strength γ_1 and γ_2 separated by a distance r, which can only take discrete values $r = \text{integer}\times b$. Therefore $2k_F r = \pi/(2b)\times\text{integer}\times b = \text{integer}\times\pi$, which means that $\exp i\chi = (-1)^{r/b}\exp i\delta$, where $\delta = -\arctan[(\gamma_1+\gamma_2)/(1-\gamma_1\gamma_2)]$. If in addition the impurities are fined tuned such that $\gamma_1\gamma_2=1$, the phase shift $\delta=-\pi/2$ and Eq. (1) becomes:

$$\begin{aligned} V_{12} &= \frac{v_F}{2\pi r}\,\Re\,\mathrm{Li}_2\left[\frac{i(-1)^{r/b}}{\gamma_1+\gamma_2}\right] \\ &= \frac{v_F}{8\pi r}\mathrm{Li}_2[-(\gamma_1+\gamma_2)^{-2}] \sim -\frac{v_F}{r}\ . \end{aligned} \tag{B1}$$

This shows that the interaction appears to be purely attractive – as $\mathrm{Li}_2(-1/4)\le \mathrm{Li}_2(-(\gamma_1+\gamma_2)^{-2})<0$ – and not oscillating, when probing inter-impurity distances r on the lattice only. This phenomenon occurs when three conditions are fulfilled: the system is half-filled ($k_F=\pi/(2b)$), the impurity separation only takes discrete values on a lattice ($r=\text{integer}\times b$) and the impurity strengths satisfy $\gamma_1\gamma_2=1$. For a numerical study of the half-filled lattice model of spinless fermions with nearest-neighbor interaction, and with two impurities, showing that this phenomenon survives in the case of interacting fermions, see Ref. [19].

---

[1] T. Giamarchi, *Quantum Physics in One Dimension* (Oxford University Press, Oxford, 2004).

[2] H. Moritz, T. Stöferle, M. Köhl, and T. Esslinger, Phys.

Rev. Lett. **91**, 250402 (2003).

[3] A. Recati, P. O. Fedichev, W. Zwerger, J. von Delft, and P. Zoller, Phys. Rev. Lett. **94**, 040404 (2005).

[4] H.W.Ch. Postma, T. Teepen, Z. Yao, M. Grifoni, and C. Dekker, Science **293**, 76 (2001).

[5] P.C. Snijders, S. Rogge, and H.H. Weitering, Phys. Rev. Lett. **96**, 076801 (2006); Europhysics News **37** (volume 5), 27-30 (2006).

[6] A. Recati, J.N. Fuchs, C.S. Peca, and W. Zwerger, Phys. Rev. A **72**, 023616 (2005).

[7] O. Kenneth, and I. Klich, Phys. Rev. Lett. **97**, 160401 (2006).

[8] A. Zee, *Quantum Field Theory in a Nutshell*, pages 65-67 (Princeton University Press, Princeton, 2004).

[9] Y. Imry, Phys. Rev. Lett. **95**, 080404 (2005).

[10] F. D. M. Haldane, Phys. Rev. Lett. **47**, 1840 (1981).

[11] L.I. Glazman, I.M. Ruzin, and B.I. Shklovskii, Phys. Rev. B **45**, 8454 (1992).

[12] H.J. Schulz, Phys. Rev. Lett. **71**, 1864 (1993).

[13] A.O. Gogolin, A.A. Nersesyan, and A.M. Tsvelik, *Bosonization and Strongly Correlated Systems* (Cambridge University Press, Cambridge, 1998), p. 298-301.

[14] C. L. Kane, and M. P. A. Fisher, Phys. Rev. Lett. **68**, 1220 (1992); Phys. Rev. B **46**, 7268 (1992); Phys. Rev. B **46**, 15233 (1992).

[15] A. Furusaki, and N. Nagaosa, Phys. Rev. B, **47**, 3827 (1993).

[16] R.L. Jaffe, Phys. Rev. D **72**, 021301 (2005).

[17] J.S. Caux, H. Saleur, and F. Siano, Phys. Rev. Lett. **88**, 106402 (2002); Nucl. Phys. B **672**, 411 (2003).

[18] See pages 318-323 in Ref. [1].

[19] P. Waechter, V. Meden, K. Schoenhammer, preprint *Indirect forces between impurities in one-dimensional quantum liquids* (unpublished). The analytical results obtained in part I and II are confirmed by the numerical calculations of Waechter *et al.*

[20] These terms, which give rise to the oscillations in the interaction energy V_{12}, were neglected in the simplified calculations of Ref. [6].

[21] This result can also be written $V_{12}/\omega_r = \pi(1-K)/24K + \Re\,\mathrm{Li}_2(-\exp i\chi)/2\pi K$.

[22] Note that $K = 1/[2(1+n)] \leq 1/4$.

[23] When $K = 1$, the solution giving a minimal free energy is $\bar{\lambda} = \max(0, 2\pi V/\omega_c - 1)$.